Author Index

Agris, P. F., see Guenther, R. H. 79

Alpert, A. J.
Hydrophobic interaction chromatography of peptides as an alternative to reversed-phase chromatography 269

Alvarez-Gonzalez, R.
Syntheisis and purification of deoxyribose analogues of NAD^+ by affinity chromatography and strong-anion-exchange high-performance liquid chromatography 89

Antle, P. E., see Cox, G. B. 325

——, see Snyder, L.R, 303

Babashak, J. V.
—— and Phillips, T. M.
Use of avidin-coated glass beads as a support for high-performance immunoaffinity chromatography 21

Barford, R. A., see Parris, N. 219

Bassoli, A.
——, Chioccara, F., Di Gregorio, G., Rindone, B., Tollari, S., Falagiani, P., Riva, G. And Bolzacchini, E.
Analysis of allergenic components of a *Parietaria judaica* pollen extract by chromatographic methods for the evaluation of purification procedures 209

Benbow, R. M., see Stowers, D. J. 47

Benedek, K.
——, Hughes, B., Seaman, M. B. and Swadesh, J. K.
Reversed-phase liquid chromatography and sodium dodecyl sulfate polyacrylamide gel electrophoresis characteristics of a recombinant DNA derived malaria antigen 191

Bolzacchini, E., see Bassoli, A. 209

Boppana, V. K., see Rhodes, G. R. 123

Borger, F. R., see Lee, J. K. 141

Burke, Jr., T. R., see Knight, M. 345

Burke, T. W. L., see Hodges, R. S. 349

Cameron Wallace, L., see Carilli, C. T. 203

Carilli, C. T.
——, Cameron Wallace, L., Smith, L. M., Wong, M. A. and Lewicki, J. A.
Semi-preparative purification of recombinant human renin and prorenin 203

Cecchini, D. J.
——, Guan, K. L. and Giese, R. W.
Staphylococcal nuclease high-performance liquid chromatographic characterization of diaminooctane-modified DNA and its biotin and fluorescein derivatives 97

Chen, F.-M.
——, Naeve, G. S. and Epstein, A. L.
Comparison of Mono Q, Superose-6, and ABx fast protein liquid chromatography for the purification of IgM monoclonal antibodies 153

Chioccara, F., see Bassoli, A. 209

Cox, G. B.
——, Antle, P. E. and Snyder, L. R.
Preparative separation of peptide and protein samples by high-performance liquid chromatography with gradient elution. II. Experimental examples compared with theory 325

——, see Snyder, L. R. 303

Davidson, C., see Lee, J. K. 141

Deluccia, F. J., see Lee, J. K. 141

Di Gregorio, F., see Bassoli, A. 209

Dutta, I.
——, O'Donovan, G. A. and Smith, D. W.
Nucleotide extraction and quantitation from tomato roots by high-performance liquid chromatography 183

Epstein, A. L., see Chen, F.-M. 153

Escribano, J.
——, Matas, R. and Méndez, E.
High-performance liquid chromatography and diode-array detection of the human protein HC (human complex-forming glycoprotein heterogeneous in charge), a chromophore-associated protein 165

Falagiani, P., see Bassoli, A. 209

Fausnaugh, J., see Tous, G. 67

Frantz, S. C., see Phillips, T. M. 13

——, Henriques, H. F. and Phillips, T. M.
Immunoaffinity of the Lewis blood group antigens 41

Ghodbane, S.
—— and Guiochon, G.
Optimization of concentration overload in preparative liquid chromatography 275

Giese, R. W., see Cecchini, D. J. 97

Gopal, D. H., see Guenther, R. H. 79

Gregorio, F. di, see Bassoli, A. 209

Gruber, H. A., see Parris, N. 219

Guan, K. L., see Cecchini, D. J. 97

Guenther, R. H.
——, Gopal, D. H. and Agris, P. F.
Purification of transfer RNA species by single-step ion-exchange high-performance liquid chromatography 79

Guiochon, G., see Ghodbane, S. 275

Habermann, R., see Josić, D. 29

Hartman, P. A.
—— and Stodola, J. D.
Reversed-phase high-performance liquid chromatography of fertirelin acetate and related compounds 177
Henriques, H. F., see Frantz, S. C. 41
Herring, S. W., see Shitanishi, K. T. 107
Hochuli, E.
Large-scale chromatography of recombinant proteins 293
Hodges, R. S.
——, Burke, T. W. L., and Mant, C. T.
Preparative purification of peptides by reversed-phase chromatography. Sample displacement mode *versus* gradient elution mode 349
Hofmann, W., see Josić, D. 29
Hudecz, F., see Szókán, G. 115
Hughes, B., see Benedek, K. 191
Hyder, S. M.
—— and Wittliff, J. L.
High-performance hydrophobic interaction chromatography as a means of identifying estrogen receptors expressing different binding domains 225
Janis, L. J.
—— and Regnier, F. E.
Immunological-chromatographic analysis 1
Josić, D.
——, and Hofmann, W., Habermann, R., and Reutter, W.
High-performance concanavalin A affinity chromatography of liver and hepatoma membrane proteins 29
Kaplan, E. D., see Siezen, R. J. 239
Keim, J. M. B., see Stowers, D. J. 47
Kelly, E. L., see Lee, J. K. 141
Knight, M.
——, Mack, G. D., Perkins, R. and Burke, Jr., T. R.
Reversed-phase preparative chromatography of [D-Ala^1]-peptide T amide 345
Lee, J. K.
——, Deluccia, F. J., Kelly, E. L., Davidson, C. and Borger, F. R.
Determination of the molecular size distribution of immunoglogin G (IgG) in intravenous IgC–albumin formulations by high-performance liquid chromatography 141
Lewicki, J. A., see Carilli, C. T. 203
Lyoo, Y. S., see Stowers, D. J. 47
Mack, G. D., see Knight,. M. 345
Mant, C. T., see Hodges, R. S. 349
Matas, R., see Escribano, J. 165
Méndez, E., see Esribano, J. 165
Merion, M., see Stowers, D. J. 47
Mezö, G., see Szókán, G. 115
Modak, M. J., see Pandey, V. N. 133
Mozersky, S. M., see Parris, N. 219
Naeve, G. S., see Chen, F.-M. 153
O'Donovan, G. A., see Dutta, I. 183
Pandey, V. N.
—— and Modak, M. J.
Purification of nucleotide-linked peptide 133
Parris, N.
——, Gruber, H. A., Mozersky, S. M. and Barford, R. A.
Purification of cytochtrome *c* from corn root mitochondria by hydrophobic interaction chromatography 219
Paul, P. S., see Stowers, D. J. 47
Perkins, R., see Knight, K.M. 345
Peterson, R. E.
—— and Wolf, W. J.
Scale-up from analytical to preparative reversed-phase high-performance liquid chromatography of soy bean glycinin 263
Phillips, T. M.
—— and Frantz, S. C.
Isolation of specific lymphocyte receptors by high-performance immunoaffinity chromatography 13
——, see Babashak, J. V. 21
——, see Frantz, S. C. 41
Regnier, F. E., see Janis, L. J. 1
Reutter, W., see Josić, D. 29
Rhodes, G. R.
—— and Boppana, V. K.
High-performance liquid chromatographic analysis of arginine-containing peptides in biological fluids by means of a selective post-column reaction with fluorescence detection 123
Rindone, B., see Bassoli, A. 209
Riva, G., see Bassoli, A. 209
Seaman, M. B., see Benedek, K. 191
Shaner, D. L., see Singh, B. K. 251
Shitanishi, K. T.
—— and Herring, S. W.
Separation of human factor X from factor Xa by reversed-phase high-performance liquid chromatography 107
Siezen, R. J.
—— and Kaplan, E. D.
Optimal resolution of eye lens γ-crystallins by cation-exchange high-performance liquid chromatography on SynChropak CM300 239
Singh, B. K.
——, Stidham, M. A. and Shaner, D. L.
Separation and characterization of two forms of acetohydroxyacid synthase from Black Mexican Sweet corn cells 251

Smith, D. W., see Dutta, I. 183
Smith, L. M., see Carilli, C. T. 203
Snyder, L. R.
——, Cox, G. B. and Antle, P. E.
Preparative separation of peptide and protein samples by high-performance liquid chromatography with gradient elution. I. The Craig model as a basis for computer simulation 303
——, see Cox, G. B. 325
Stein, S., see Tous, G. 67
Stidham, M. A., see Singh, B. K. 251
Stodola, P. A., see Hartman, P. A. 177
Stowers, D. J.
——, Keim, J. M. B., Paul, P. S., Lyoo, Y. S., Merion, M. and Benbow, R. M.
High-resolution chromatography of nucleic acids on the Gen-Pak PAX column 47
Swadesh, J. K., see Benedek, K. 191
Szókán, G.
——, Mezö, G. and Hudecz, F.
Application of Marfey's reagent in racemization studies of amino acids and peptides 115
Tollari, S., see Bassoli, A. 209
Tous, G.
——, Fausnaugh, J., Vieira, P. and Stein, S.
Preparation and chromatographic use of 5′-fluorescent-labelled DNA probes 67
Vieira, P., see Tous, G. 67
Wittliff, J. L., see Hyder, S. M. 225
Wolf, W. J., see Peterson, R. E. 263
Wong, M. A., see Carilli, C. T. 203